arXiv:0903.5138v1 [hep-ph] 30 Mar 2009

Radiative decay into γ-baryon decuplet of dynamically generated resonances

Bao-Xi Sun[1] and E. Oset[2]

March 30, 2009

[1] Institute of Theoretical Physics, College of Applied Sciences,
Beijing University of Technology, Beijing 100124, China
[2] Departamento de Física Teórica and IFIC, Centro Mixto Universidad de Valencia-CSIC,
Institutos de Investigación de Paterna, Aptdo. 22085, 46071 Valencia, Spain

Abstract

We study the radiative decay into γ and a baryon of the SU(3) decuplet of ten resonances that are dynamically generated from the interaction of vector mesons with baryons of the decuplet. We obtain quite different partial decay widths for the various resonances, and for different charge states of the same resonance, suggesting that the experimental investigation of these radiative decays should bring much information on the nature of these resonances.

1 Introduction

In a recent paper [1], the $\rho\Delta$ interaction was studied within the local hidden gauge formalism for the interaction of vector mesons. The results of the interaction gave a natural interpretation for the $\Delta(1930)(5/2^-)$ as a $\rho\Delta$ bound state, which otherwise is extremely problematic in quark models since it involves a $3h\omega$ excitation and appears with much higher mass. At the same time two states with $J^P = 1/2^-, 3/2^-$ were obtained, degenerate with the $5/2^-$, which could be accommodated with two more known Δ states in that energy range. Also, three degenerate N^* states with $1/2^-, 3/2^-, 5/2^-$ were obtained, which were more difficult to identify with known resonances since that sector is not so well established. The work of [1] was extended to the SU(3) sector in [2] to account for the interaction of vectors of the octet with baryons of the decuplet. In this case ten resonances, all of them also degenerate in the three spin states, were obtained, many of which could be identified with existing resonances, while there were predictions for a few more. At the same time in [2] the poles and residues at the poles of the resonances were evaluated, providing the coupling of the resonances to the different vector-baryon of the decuplet components.

One of the straightforward tests of these theoretical predictions is the radiative decay of these resonances into photon and the member of the baryon decuplet to which it couples. Radiative decay of resonances into γN is one of the observables traditionally calculated in hadronic models. Work in quark models on this issue is abundant [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18]. For resonances which appear as dynamically generated in chiral unitary theories there is also much work done on the radiative decay into γN [19, 20, 21, 22, 23]. Experimental work in this topic is also of current interest [25, 26, 27].

In the present work we address the novel aspect of radiative decay into a photon and a baryon of the decuplet of the Δ, since the undelying dynamics of the resonances that we study provides this as the dominant mode of radiative decay into photon baryon. This is so, because the underlying theory of the studies of [1, 2] is the local hidden gauge formalism for the interaction of vector mesons developed in [28, 29, 30], which has the peculiar feature, inherent to vector meson dominance, that the photons couple to the hadrons through the conversion into a vector meson. In this case a photon in the final state comes from either a ρ^0, ω, ϕ. Thus, the radiative decay of the resonances into γB is readily obtained from the theory by taking the terms with $\rho^0 B, \omega B, \phi B$ in the final state and coupling the γ to any of the final ρ^0, ω, ϕ vector mesons. This procedure was used in [31] and provided good results for the radiative decay into $\gamma\gamma$ of the $f_0(1370)$ and $f_2(1270)$ mesons which were dynamically generated from the $\rho\rho$ interaction within the same framework [32]. This latter work was also extended to the interaction of vectors with themselves within SU(3), where many other states are obtained which can be also associated with known resonances [33]. Given the success of the theory in its predictions and the good results obtained for the $\gamma\gamma$ decay of the $f_0(1370)$ and $f_2(1270)$ mesons, the theoretical framework stands on good foot and the predictions made should be solid enough to constitute a test of the theory by contrasting with experimental data.

The experimental situation in that region of energies is still poor. The PDG [34] quotes many radiative decays of N^* resonances, and of the $A_{1/2}, A_{3/2}$ helicity amplitudes for decay of resonances into γN, with N either proton or neutron. However, there are no data to our knowledge for radiative decay into γB, with B a baryon of the decuplet. The reason for it might be the difficulty in the measurement, or the lack of motivation, since there are also no theoretical works devoted to the subject. With the present work we hope to reverse the situation offering a clear motivation for these experiments since they bear close connection with the nature invoked for these resonances, very different to the ordinary three quark structure of the baryons. In fact, the numbers obtained for the radiative widths are well within measurable range, of the order of 1 MeV, and the predictions are interesting, with striking differences of one order of magnitude between decay widths for different charges of the same resonance.

The work will proceed as follows. In the next two Sections we present the framework for the evaluation of amplitudes of radiative decay. In Section 4 we show the results obtained for the different resonances and in Section 5 we finish with some conclusions.

2 Framework

In Ref. [1, 2], the scattering amplitudes for vector-decuplet baryon $VB \to V'B'$ are given by

$$t_{VB\to V'B'} \;=\; t\;\vec{\epsilon}\cdot\vec{\epsilon'}\;\delta_{m_s,m'_s}, \tag{1}$$

where $\vec{\epsilon}$, $\vec{\epsilon'}$ refer to the initial and final vector polarization and the matrix is diagonal in the third component of the baryons of the decuplet. The transition is diagonal in spin of the baryon and spin of the vector, and as a consequence in the total spin. To make this property more explicit, we write the states of total spin as

$$|S,M\rangle \;=\; \sum_{m_s} C\,(3/2,1,S;m_s,M-m_s,M)\,|3/2,m_s\rangle|\vec{\epsilon}_{M-m_s}\rangle \tag{2}$$

and

$$\langle S,M| \;=\; \sum_{m'_s} C\,(3/2,1,S;m'_s,M-m'_s,M)\,\langle 3/2,m'_s|\langle\vec{\epsilon}^{\,*}_{M-m'_s}|, \tag{3}$$

where $C\,(3/2,1,S;m_s,M-m_s,M)$ are the Clebsch-Gordan coefficients and ϵ_μ the polarization vectors in spherical basis

$$\vec{\epsilon}_+ \;=\; -\frac{1}{\sqrt{2}}\left(\vec{\epsilon}_1 \,+\, i\vec{\epsilon}_2\right), \quad \vec{\epsilon}_- \;=\; \frac{1}{\sqrt{2}}\left(\vec{\epsilon}_1 \,-\, i\vec{\epsilon}_2\right), \quad \vec{\epsilon}_0 \;=\; \vec{\epsilon}_3. \tag{4}$$

We can write Eq. (1) in terms of the projectors $|S,M\rangle\langle S,M|$ as

$$t_{VB\to V'B'} \;=\; t\;\langle\vec{\epsilon'}|\langle 3/2,m'_s|\sum_{S,M}|S,M\rangle\langle S,M|3/2,m_s\rangle|\vec{\epsilon}\,\rangle. \tag{5}$$

Since the Clebsch-Gordan coefficients satisfy the normalization condition

$$\sum_S C\,(3/2,1,S;m_s,M-m_s,M)\,C\,(3/2,1,S;m'_s,M'-m'_s,M') \;=\; \delta_{m_s m'_s}\delta_{MM'}, \tag{6}$$

we have then

$$\begin{aligned}\sum_{S,M}|S,M\rangle\langle S,M| \;&=\; \sum_M\sum_{m_s}|3/2,m_s\rangle\langle 3/2,m_s|\;|\vec{\epsilon}_{M-m_s}\;\rangle\langle\vec{\epsilon^*}_{M-m_s}| \\ &=\; \sum_{M'}\sum_{m_s}|3/2,m_s\rangle\langle 3/2,m_s|\;|\vec{\epsilon}_{M'}\;\rangle\langle\vec{\epsilon^*}_{M'}| \equiv 1.\end{aligned} \tag{7}$$

We can depict the contribution of a specific resonant state of spin S to the amplitude described by means of Fig. 1. Then the amplitude for the transition of the resonance to a final vector-baryon state is depicted by means of Fig. 2. As shown is Ref. [1, 2], the $VB \to V'B'$ scattering amplitudes develop poles corresponding to resonances and a resonant amplitude is written as Eq. (1) with t given by

$$t_{ij} \;=\; \frac{g_i g_j}{\sqrt{s}-M+i\Gamma/2} \tag{8}$$



Figure 1: Diagram contributing to the vector-baryon interaction via the exchange of a resonance.



Figure 2: Diagram on the decay of the resonance in a decuplet baryon and a vector meson.

with g_i and g_j the couplings to the initial and final states. Accordingly, the amplitude for the transition from the resonance to a final state of vector-baryon is given by

$$\begin{aligned} t_{SM\rightarrow V'B'} &= g_i\langle\vec{\epsilon}|\langle 3/2, m_s|S, M\rangle \\ &= g_i C(3/2, 1, S; m_s, M-m_s, M)\langle\vec{\epsilon}\,|\vec{\epsilon}_{M-m_s}\rangle. \end{aligned} \tag{9}$$

When calculating the decay width of the resonance into VB we will sum $|t|^2$ over the vector and baryon polarization, and average over the resonance polarization M. Thus, we have

$$\begin{aligned} &\frac{1}{2S+1}\sum_{M,m_s,\vec{\epsilon}}|t_{SM\rightarrow V'B'}|^2 \qquad (10)\\ =\ & |g_i|^2\frac{1}{2S+1}\sum_{M,m_s,\vec{\epsilon}} C(3/2,1,S;m_s,M-m_s,M)^2\langle\vec{\epsilon}^{\,*}{}_{M-m_s}|\vec{\epsilon}\rangle\langle\vec{\epsilon}|\vec{\epsilon}_{M-m_s}\rangle \\ =\ & |g_i|^2\frac{1}{2S+1}\sum_{M'}\sum_{m_s}\frac{2S+1}{3}C(3/2,S,1;m_s,m_s+M',-M')^2\langle\vec{\epsilon}^{\,*}{}_{M'}|\vec{\epsilon}_{M'}\rangle \\ =\ & |g_i|^2\frac{1}{3}\sum_{M'}\delta_{M'M'} \\ =\ & |g_i|^2, \end{aligned}$$

where in the first step we have permuted the two last spins in the Clebsch-Gordan coefficients and in the second we applied their orthogonality condition.

We observe that the normalization of the amplitudes is done in a way such that the sum and average of $|t|^2$ is simply the modulus squared of the coupling of the resonance to the final state. The width of the resonance for decay into VB is given by

$$\Gamma \;=\; \frac{M_B}{2\pi M_R}\, q\, |g_i|^2, \tag{11}$$

where q is the momentum of the vector in the resonance rest frame and M_B, M_R the masses of the baryon and the resonance. We should note already that later on when the vector polarizations are substituted by the photon polarizations in the sum over M' in Eq. (10) we will get a factor two rather than three, because we only have two transverse polarizations, and then Eq. (11) must be multiplied by the factor 2/3.

3 Radiative decay

Next we study the radiative decay into $B\gamma$ of the resonances dynamically generated in Ref. [2] with B a baryon of the decuplet. Recalling the results of [2] we obtained there ten resonances dynamically generated, each of them degenerated in three states of spin, $1/2^-$, $3/2^-$, $5/2^-$. As we have discussed in the former section, the radiative width will not depend on the spin of the resonance, but only on the coupling which is the same for all three spin states due to the degeneracy. This would be of course an interesting experimental test of the nature of these resonances.



Figure 3: Diagram on the radiative decay of the resonance in a decuplet baryon and a photon.

In order to proceed further, we use the same formalism of the hidden gauge local symmetry for the vector mesons of [28, 29, 30]. The peculiarity of this theory concerning photons is that they couple to hadrons by converting first into a vector meson, ρ^0, ω, ϕ. Diagrammatically this is depicted in Fig. 3. This idea has already been applied with success to obtain the radiative decay of the $f_0(1270)$ and $f_0(1370)$ resonances into $\gamma\gamma$ in Ref. [31]. In that work the question of gauge invariance was addressed and it was shown that the theory fulfills it. In Ref. [35], it is also proved in the case of radiative decay of axial vector resonances.

The amplitude of Fig. 3 requires the γV convertion Lagrangian, which comes from Refs. [28, 29, 30] and is given by (*see Ref.* [35] *for practical details*)

$$\mathcal{L}_{V\gamma} \;=\; -M_V^2 \frac{e}{\tilde{g}} A_\mu \langle V^\mu Q\rangle \tag{12}$$

with A_μ the photon field, V_μ the SU(3) matrix of vector fields

$$V_\mu \equiv \left(\begin{array}{ccc} \frac{1}{\sqrt{2}}\rho^0 + \frac{1}{\sqrt{2}}\omega & \rho^+ & K^{*+} \\ \rho^- & -\frac{1}{\sqrt{2}}\rho^0 + \frac{1}{\sqrt{2}}\omega & K^{*0} \\ K^{*-} & \bar{K}^{*0} & \phi \end{array}\right)_\mu , \tag{13}$$

and Q the charge matrix

$$Q \equiv \left(\begin{array}{ccc} 2/3 & 0 & 0 \\ 0 & -1/3 & 0 \\ 0 & 0 & -1/3 \end{array}\right). \tag{14}$$

In Eq. (12), M_V is the vector meson mass, for which we take an average value $M_V = 800 MeV$, e the electron charge, $e^2 = 4\pi\alpha$, and

$$\tilde{g} = \frac{M_V}{2f}; \qquad f = 93 MeV.$$

The sum over polarizations in the intermediate vector meson, which converts the polarization vector of the vector meson of the $R \to BV$ amplitude into the photon polarization of the $R \to B\gamma$ amplitude, leads to the equation

$$-it_{\gamma V} D_V \;=\; -iM_V^2 \,\frac{e}{\tilde{g}}\, \frac{i}{-M_V^2}\, F_j \tag{15}$$

with

$$F_j \;=\; \left\{ \begin{array}{ll} \frac{1}{\sqrt{2}} & for \;\; \rho^0, \\ \frac{1}{3\sqrt{2}} & for \;\; \omega, \\ -\frac{1}{3} & for \;\; \phi. \end{array}\right. \tag{16}$$

Thus, finally our amplitude for the $R \to B\gamma$ transition, omitting the spin matrix element of Eq. (9), is given by

$$t_\gamma \;=\; -\frac{e}{\tilde{g}} \sum_{j=\rho^0,\ \omega,\ \phi} g_j F_j. \tag{17}$$

As discussed in the former section, the radiative decay width will then be given by

$$\Gamma_\gamma \;=\; \frac{1}{2\pi}\,\frac{2}{3}\,\frac{M_B}{M_R}\, q\, |t_\gamma|^2. \tag{18}$$

The couplings g_j for different resonance and VB with $V = \rho^0,\ \omega,\ \phi$ and B different baryon of the decuplet can be found in Ref. [2] and we use them here for the evaluation of Γ_γ. The factor $\frac{2}{3}$ in eq. (18) additional to eq. (11) appears because now we have only two photon polarizations and the sum over M' in eq. (10) gives 2 instead of 3 for the case of vector mesons.

4 Results

The couplings of the resonances to the different VB channels are given in Ref. [2] in the isospin basis. For the case of ωB and ϕB, there is no change to be done, but for the case of ρB, one must project over the $\rho^0 B$ component. Since this depends on the charge of the resonance R, the radiative decays will depend on this charge, as we will see. We recall that in our phase convention $|\rho^+\rangle = -|1,1\rangle$ of isospin. The information on the resonances and their couplings to different baryons of decuplet and vector mesons ρ,ω, ϕ for different channels is listed in Table 1. We detail the results below.

S, I	Channel			
		$z_R = 1850 + i5$		
0, 1/2	$\Delta\rho$	$4.9 + i0.1$		
		$z_R = 1972 + i49$		
0, 3/2	$\Delta\rho$ $\Delta\omega$ $\Delta\phi$	$5.0 + i0.2$ $-0.1 + i0.2$ $0.2 - i0.4$		
		$z_R = 2052 + i10$		
-1, 0	$\Sigma^*\rho$	$4.2 + i0.1$		
		$z_R = 1987 + i1$	$z_R = 2145 + i58$	$z_R = 2383 + i73$
-1,1	$\Sigma^*\rho$ $\Sigma^*\omega$ $\Sigma^*\phi$	$1.4 + i0.0$ $1.4 + i0.0$ $-2.1 - i0.0$	$-4.3 - i0.7$ $1.3 - i0.4$ $-1.9 + i0.6$	$0.4 + i1.1$ $-1.4 - i0.4$ $2.1 + i0.6$
		$z_R = 2214 + i4$	$z_R = 2305 + i66$	$z_R = 2522 + i38$
-2, 1/2	$\Xi^*\rho$ $\Xi^*\omega$ $\Xi^*\phi$	$1.8 - i0.1$ $1.7 + i0.1$ $-2.5 - i0.1$	$-3.5 - i1.7$ $2.0 - i0.7$ $-3.0 + i1.0$	$0.2 + i1.0$ $-0.6 - i0.3$ $0.9 + i0.4$
		$z_R = 2449 + i7$		
-3, 0	$\Omega\omega$ $\Omega\phi$	$1.6 - i0.2$ $-2.4 + i0.3$		

Table 1: The coupling g_i of the resonance obtained dynamically to the ρB, ωB and ϕB channels.

4.1 $S = 0, I = 1/2$ channel

A resonance is obtained at $z_R = 1850 + i5 MeV$ which couples to $\Delta\rho$. We have in this case

$$|\Delta\rho, \frac{1}{2}, \frac{1}{2}\rangle \ = \ \sqrt{\frac{1}{2}}|\Delta^{++}\rho^-\rangle \ - \ \sqrt{\frac{1}{3}}|\Delta^+\rho^0\rangle \ - \ \sqrt{\frac{1}{6}}|\Delta^0\rho^+\rangle \tag{19}$$

and

$$|\Delta\rho, \frac{1}{2}, -\frac{1}{2}\rangle \;=\; \sqrt{\frac{1}{6}}|\Delta^{+}\rho^{-}\rangle \;-\; \sqrt{\frac{1}{3}}|\Delta^{0}\rho^{0}\rangle \;-\; \sqrt{\frac{1}{2}}|\Delta^{-}\rho^{+}\rangle, \tag{20}$$

The coupling of the resonance to ρ^0 is obtained multiplying the coupling of Table 1 by the corresponding Clebsch-Gordan coefficient for $\Delta\rho^0$ of Eqs. (19, 20). Then by means of Eqs. (17, 18), one obtains the decay width. In this case since the $\Delta\rho^0$ component is the same for $I_3 = 1/2$ and $I_3 = -1/2$, one obtains the same radiative width for the two channels, which is $\Gamma = 0.722 MeV$.

4.2 $S = 0, I = 3/2$ channel

One resonance is obtained at $z_R = 1972 + i49 MeV$ which couples to $\Delta\rho$, $\Delta\omega$ and $\Delta\phi$. The isospin states for $\Delta\rho$ can be written as

$$|\Delta\rho, \frac{3}{2}, \frac{3}{2}\rangle \;=\; \sqrt{\frac{3}{5}}|\Delta^{++}\rho^{0}\rangle \;+\; \sqrt{\frac{2}{5}}|\Delta^{+}\rho^{+}\rangle, \tag{21}$$

$$|\Delta\rho, \frac{3}{2}, \frac{1}{2}\rangle \;=\; \sqrt{\frac{2}{5}}|\Delta^{++}\rho^{-}\rangle \;+\; \sqrt{\frac{1}{15}}|\Delta^{+}\rho^{0}\rangle \;+\; \sqrt{\frac{8}{15}}|\Delta^{0}\rho^{+}\rangle, \tag{22}$$

$$|\Delta\rho, \frac{3}{2}, -\frac{1}{2}\rangle \;=\; \sqrt{\frac{8}{15}}|\Delta^{+}\rho^{-}\rangle \;-\; \sqrt{\frac{1}{15}}|\Delta^{0}\rho^{0}\rangle \;+\; \sqrt{\frac{2}{5}}|\Delta^{-}\rho^{+}\rangle, \tag{23}$$

$$|\Delta\rho, \frac{3}{2}, -\frac{3}{2}\rangle \;=\; \sqrt{\frac{2}{5}}|\Delta^{0}\rho^{-}\rangle \;-\; \sqrt{\frac{3}{5}}|\Delta^{-}\rho^{0}\rangle. \tag{24}$$

Since all the Clebsch-Gordan coefficients to $\Delta\rho^0$ are now different, we obtain different radiative decay width for each charge of the state. The results are $\Gamma = 1.402 MeV$ for $I_3 = 3/2$, $\Gamma = 0.143 MeV$ for $I_3 = 1/2$, $\Gamma = 0.203 MeV$ for $I_3 = -1/2$ and $\Gamma = 1.582 MeV$ for $I_3 = -3/2$. It is quite interesting to see that there is an order of magnitude difference between for $I = 3/2$ and $I = 1/2$, and it is a clear prediction that could be tested experimentally.

4.3 $S = -1, I = 0$ channel

We get a resonance at $z_R = 2052 + i10 MeV$, which couples to $\Sigma^*\rho$. In this case

$$|\Sigma^*\rho, 0, 0\rangle \;=\; \sqrt{\frac{1}{3}}|\Sigma^{*+}\rho^{-}\rangle \;-\; \sqrt{\frac{1}{3}}|\Sigma^{*0}\rho^{0}\rangle \;-\; \sqrt{\frac{1}{3}}|\Sigma^{*-}\rho^{+}\rangle, \tag{25}$$

and the radiative decay obtained is $\Gamma \;=\; 0.583 MeV$.

4.4 $S = -1, I = 1$ channel

Here we find three resonances at $z_R = 1987 + i1MeV$, $2145 + i58MeV$ and $2383 + i73MeV$, which couple to $\Sigma^*\rho$, $\Sigma^*\omega$ and $\Sigma^*\phi$. The relevant isospin states are

$$|\Sigma^*\rho, 1, 1\rangle = \sqrt{\frac{1}{2}}|\Sigma^{*+}\rho^0\rangle + \sqrt{\frac{1}{2}}|\Sigma^{*0}\rho^+\rangle, \tag{26}$$

$$|\Sigma^*\rho, 1, 0\rangle = \sqrt{\frac{1}{2}}|\Sigma^{*+}\rho^-\rangle + \sqrt{\frac{1}{2}}|\Sigma^{*-}\rho^+\rangle, \tag{27}$$

and

$$|\Sigma^*\rho, 1, -1\rangle = \sqrt{\frac{1}{2}}|\Sigma^{*0}\rho^-\rangle - \sqrt{\frac{1}{2}}|\Sigma^{*-}\rho^0\rangle. \tag{28}$$

The results obtained in this case are summarized in Table 2.

I_3	(1987)	(2145)	(2383)
1	0.561	0.399	0.182
0	0.199	0.206	0.277
−1	0.020	2.029	0.537

Table 2: The radiative decay widths in units of MeV for the $S = -1, I = 1$ resonances with different isospin projection I_3.

4.5 $S = -2, I = \frac{1}{2}$ channel

Here we also find three states at $z_R = 2214 + i4MeV$, $2305 + i66MeV$ and $2522 + i38MeV$, which couple to $\Xi^*\rho$, $\Xi^*\omega$ and $\Xi^*\phi$. The isospin states for $\Xi^*\rho$ are written as

$$|\Xi^*\rho, \frac{1}{2}, \frac{1}{2}\rangle = \sqrt{\frac{2}{3}}|\Xi^{*-}\rho^+\rangle + \sqrt{\frac{1}{3}}|\Xi^{*0}\rho^0\rangle, \tag{29}$$

$$|\Xi^*\rho, \frac{1}{2}, -\frac{1}{2}\rangle = -\sqrt{\frac{1}{3}}|\Xi^{*-}\rho^0\rangle + \sqrt{\frac{2}{3}}|\Xi^{*0}\rho^-\rangle, \tag{30}$$

The radiative decay widths in this case are shown in Table 3.

4.6 $S = -3, I = 0$ channel

Here we have only one state at $z_R = 2449 + i7MeV$, which couples to $\Omega\omega$ and $\Omega\phi$. The radiative decay width obtained in this case is $\Gamma = 0.330MeV$.

As one can see, there is a large variation in the radiative width of the different states, which should constitute a good test for the model when these widths are measured.

In Table 4 we summarize all the results obtained making an association of our states to some resonances found in the PDG[34].

I_3	(2214)	(2305)	(2522)
1/2	0.815	0.320	0.044
−1/2	0.054	1.902	0.165

Table 3: The radiative decay widths in units of MeV for the $S = -2, I = 1/2$ resonances with the different isospin projection I_3.

S, I	Theory	PDG data		Predicted width (KeV) for I_3						
	pole position (MeV)	name	J^P	−3/2	−1	−1/2	0	1/2	1	3/2
$0, 1/2$	$1850 + i5$	$N(2090)$	$1/2^-$			722		722		
		$N(2080)$	$3/2^-$							
$0, 3/2$	$1972 + i49$	$\Delta(1900)$	$1/2^-$	1582		203		143		1402
		$\Delta(1940)$	$3/2^-$							
		$\Delta(1930)$	$5/2^-$							
$-1, 0$	$2052 + i10$	$\Lambda(2000)$	$?^?$				583			
$-1, 1$	$1987 + i1$	$\Sigma(1940)$	$3/2^-$		20		199		561	
	$2145 + i58$	$\Sigma(2000)$	$1/2^-$		2029		206		399	
	$2383 + i73$	$\Sigma(2250)$	$?^?$		537		277		182	
		$\Sigma(2455)$	$?^?$							
$-2, 1/2$	$2214 + i4$	$\Xi(2250)$	$?^?$			54		815		
	$2305 + i66$	$\Xi(2370)$	$?^?$			1902		320		
	$2522 + i38$	$\Xi(2500)$	$?^?$			165		44		
$-3, 1$	$2449 + i7$	$\Omega(2470)$	$?^?$				330			

Table 4: The predicted radiative decay widths of the 10 dynamically generated resonances for different isospin projection I_3. Their possible PDG counterparts are also listed. Note that the $\Sigma(2000)$ could be the spin parter of the $\Sigma(1940)$, in which case the radiative decay widths would be those of the $\Sigma(1940)$.

5 Conclusions

We have studied the radiative decay into γB, with B a baryon of the decuplet of SU(3), of the dynamically generated resonances obtained within the framework of the local hidden gauge mechanism for vector interactions. The framework is particularly rewarding for the study of such observable, since the photon in the final state appears coupling directly to the vector $V = \rho^0, \omega, \phi$ in the $R \to V$ amplitudes which are studied in previous works. The rates obtained are large and the radiative widths are of the order of 1 MeV. On the other hand, one of the appealing features of the results is the large difference, of about one order of magnitude, that one finds between the widths for different charge states of the same particle. These results are tied to details of the theory, concretely the coupling of the resonance to $V\ B$, which sometimes produce large interferences between the different contributions of the three vector mesons to which the photon couples. As a consequence the radiative decay widths that we have evaluated bare much information on the nature of those resonances, which should justify efforts for a systematic measurement of these observables. We hope the present work stimulates work in this direction.

Acknowledgments

We would like to thank M. J. Vicente Vacas for a critical reading of the manuscript. This work is partly supported by DGICYT contract number FIS2006-03438. This research is part of the EU Integrated Infrastructure Initiative Hadron Physics Project under contract number RII3-CT-2004-506078. B. X. Sun acknowledges support from the National Natural Science Foundation of China under grant number 10775012.

References

[1] P. Gonzalez, E. Oset and J. Vijande, Phys. Rev. C **79**, 025209 (2009) [arXiv:0812.3368 [hep-ph]].

[2] S. Sarkar, B. X. Sun, E. Oset and M. J. V. Vacas, arXiv:0902.3150 [hep-ph].

[3] J. W. Darewych, M. Horbatsch and R. Koniuk, Phys. Rev. D **28**, 1125 (1983).

[4] M. Warns, W. Pfeil and H. Rollnik, Phys. Lett. B **258**, 431 (1991).

[5] Y. Umino and F. Myhrer, Nucl. Phys. A **529** (1991) 713.

[6] Y. Umino and F. Myhrer, Nucl. Phys. A **554**, 593 (1993) [arXiv:nucl-th/9210018].

[7] R. Bijker, F. Iachello and A. Leviatan, Annals Phys. **284**, 89 (2000) [arXiv:nucl-th/0004034].

[8] F. E. Close and Z. P. Li, Phys. Rev. D **42** (1990) 2194.

[9] W. Konen and H. J. Weber, Phys. Rev. D **41**, 2201 (1990).

[10] M. Warns, H. Schroder, W. Pfeil and H. Rollnik, Z. Phys. C **45**, 627 (1990).

[11] S. Capstick and B. D. Keister, Phys. Rev. D **51** (1995) 3598

[12] B.D. Keister, S. Capstick, N* Physics, edited by T.-S. H. Lee, W. Roberts (World Scientific, Singapore, 1997) p. 58.

[13] M. Aiello, M. M. Giannini and E. Santopinto, J. Phys. G **24**, 753 (1998)

[14] E. Pace, G. Salme and S. Simula, Few Body Syst. Suppl. **10**, 407 (1999)

[15] D. Merten, U. Loring, K. Kretzschmar, B. Metsch and H. R. Petry, Eur. Phys. J. A **14** (2002) 477

[16] M. M. Giannini, E. Santopinto and A. Vassallo, Prog. Part. Nucl. Phys. **50** (2003) 263 Annals Phys. **284**, 89 (2000) [arXiv:nucl-th/0004034].

[17] T. Van Cauteren, D. Merten, T. Corthals, S. Janssen, B. Metsch, H. R. Petry and J. Ryckebusch, Eur. Phys. J. A **20**, 283 (2004) [arXiv:nucl-th/0310058].

[18] T. Van Cauteren, J. Ryckebusch, B. Metsch and H. R. Petry, Eur. Phys. J. A **26**, 339 (2005) [arXiv:nucl-th/0509047].

[19] N. Kaiser, T. Waas and W. Weise, Nucl. Phys. A **612**, 297 (1997).

[20] B. Borasoy, E. Marco and S. Wetzel, Phys. Rev. C **66**, 055208 (2002)

[21] M. Döring, E. Oset and S. Sarkar, Phys. Rev. C **74** (2006) 065204

[22] M. Döring, Nucl. Phys. A **786**, 164 (2007)

[23] L. S. Geng, E. Oset and M. Döring, Eur. Phys. J. A **32**, 201 (2007)

[24] D. Jido, M. Doering and E. Oset, Phys. Rev. C **77**, 065207 (2008) [arXiv:0712.0038 [nucl-th]].

[25] R. Thompson *et al.* [CLAS Collaboration], Phys. Rev. Lett. **86**, 1702 (2001)

[26] V. D. Burkert and T. S. H. Lee, Int. J. Mod. Phys. E **13** (2004) 1035

[27] I. G. Aznauryan, V. D. Burkert, H. Egiyan, K. Joo, R. Minehart and L. C. Smith, Phys. Rev. C **71**, 015201 (2005)

[28] M. Bando, T. Kugo, S. Uehara, K. Yamawaki and T. Yanagida, Phys. Rev. Lett. **54**, 1215 (1985).

[29] M. Bando, T. Kugo and K. Yamawaki, Phys. Rept. **164**, 217 (1988).

[30] M. Harada and K. Yamawaki, Phys. Rept. **381**, 1 (2003) [arXiv:hep-ph/0302103].

[31] H. Nagahiro, J. Yamagata-Sekihara, E. Oset and S. Hirenzaki, arXiv:0809.3717 [hep-ph].

[32] R. Molina, D. Nicmorus and E. Oset, Phys. Rev. D **78**, 114018 (2008) [arXiv:0809.2233 [hep-ph]].

[33] L. S. Geng and E. Oset, Phys. Rev. D in print, arXiv:0812.1199 [hep-ph].

[34] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).

[35] H. Nagahiro, L. Roca, A. Hosaka and E. Oset, Phys. Rev. D **79**, 014015 (2009) [arXiv:0809.0943 [hep-ph]].